

## ***Lyme Disease***


### **Introduction**

Lyme disease is an infectious disease. Lyme disease is spread by a deer tick. If not treated, Lyme disease is a potentially serious disease that can lead to disabling arthritis and neurological problems.

This reference summary will help you understand Lyme disease and how it can be treated.

### **What is Lyme Disease?**

In the early 1970s, several children in Lyme, Connecticut, and in surrounding areas, developed arthritis. Health care providers soon found out that the arthritis was caused by an infectious disease, which they called Lyme disease. Lyme disease is caused by bacteria called *Borrelia burgdorferi*.



The percentage of deer ticks infected varies from region to region, reaching 50% in some areas. The national average is about 1%. These ticks are called deer ticks because they usually feed and mate on deer. When a deer tick bites a person, the bacteria can transfer to the person's blood. The tick must stay attached to the person's skin for 2 days for infection to occur.

A person bitten by an infected deer tick develops a rash called erythema migrans or EM rash. It is usually the first symptom. The rash starts as a small red spot at the site of the tick bite. The spot expands over a period of days or weeks, forming a circle or oval shaped rash.



EM Rash

Over time, the bacteria travel through the blood stream and lymph nodes. They divide and multiply and produce harmful substances called toxins. Lyme disease is not contagious.

---

This document is for informational purposes and is not intended to be a substitute for the advice of a doctor or healthcare professional or a recommendation for any particular treatment plan. Like any printed material, it may become out of date over time. It is important that you rely on the advice of a doctor or a healthcare professional for your specific condition.

## Symptoms

An EM rash is similar to a target; it looks like a red ring around a clear area with a red center. The rash can range from the size of a dime to the width of a person's back. Appearing within a few weeks, the infection spreads and the rash may appear in different places on the body.

An EM rash is often accompanied by the following symptoms:

- Fever.
- Headache.
- Stiff neck.
- Body aches.
- Fatigue.

Although Lyme disease symptoms are flu-like, they usually do not go away like the common flu. They may also occur off and on continually.



Arthritis usually shows up after several months of infection. More than half of Lyme disease patients who do not get antibiotic treatment have repeated attacks of painful, swollen joints. The arthritis may shift from one joint to another. But the knee is most commonly affected. About 10 to 20% of untreated Lyme disease patients develop chronic arthritis, which lasts a lifetime.

Lyme disease can also affect the nervous system, causing:

- Stiff neck.
- Severe headache.
- Temporary paralysis of facial muscles.
- Numbness, pain or weakness in the limbs.
- Poor motor coordination.

Other symptoms such as memory loss, difficulty concentrating and change in mood or sleeping habits are also associated with Lyme disease. Problems with the nervous system may develop several weeks, months or even years, after an untreated infection. These symptoms last for weeks or months and may recur.

---

This document is for informational purposes and is not intended to be a substitute for the advice of a doctor or healthcare professional or a recommendation for any particular treatment plan. Like any printed material, it may become out of date over time. It is important that you rely on the advice of a doctor or a healthcare professional for your specific condition.

Less than 10% of Lyme disease patients develop heart problems. Symptoms include irregular heartbeat, dizziness and shortness of breath. These symptoms usually show up a few weeks after infection and only last a few days or weeks.

Lyme disease can result in symptoms such as eye inflammation, hepatitis and severe fatigue. None of these problems are likely to appear without one or more other Lyme disease symptoms.

## Diagnosis

The most obvious symptom of Lyme disease is an EM rash. But about 25% of people who get infected do not develop an EM rash. Without an EM rash, Lyme disease can be difficult to diagnose since many of the symptoms are also the symptoms of other diseases.

Many Lyme disease patients cannot remember being bitten by a tick. This is not surprising, since the tick is tiny and its bite is usually painless. To diagnose Lyme disease, the health care provider asks the patient questions about recent activities and unusual signs he or she may have noticed. This process is called taking a medical history.

When a patient with possible Lyme disease symptoms does not develop an EM rash, the health care provider needs a blood test to check for Lyme disease.

The health care provider may request a spinal tap for patients who have nervous system symptoms. During a spinal tap, fluid is withdrawn from around the brain and spinal cord. The fluid is tested for Lyme antibodies.

## Treatment

Most Lyme disease patients can be successfully treated with antibiotics. In general, the sooner such treatment is given, the quicker and more complete the recovery is. Taking antibiotics by mouth for a few weeks can speed up healing of the EM rash and usually prevents symptoms such as arthritis or neurological problems. With more advanced or serious infection, antibiotics administered directly into the blood through an intravenous line may be needed.



---

This document is for informational purposes and is not intended to be a substitute for the advice of a doctor or healthcare professional or a recommendation for any particular treatment plan. Like any printed material, it may become out of date over time. It is important that you rely on the advice of a doctor or a healthcare professional for your specific condition.

Along with antibiotics, Lyme arthritis may also be treated with anti-inflammatory drugs. Fluid can be drawn from affected joints. The lining of inflamed joints may also be removed surgically. Sometimes it takes years for Lyme arthritis to fade away completely, but it usually goes away within a few weeks or months of antibiotic therapy.

Unfortunately, being infected with Lyme disease once does not guarantee that it will never happen again. Lyme disease can affect the same person more than once if he or she is re-infected with the Lyme disease bacterium.

## Prevention

You can do several things to avoid being exposed to deer ticks.

- Avoid tick-infested areas, especially in May, June and July.
- Wear long-sleeved shirts, socks and closed shoes when outdoors.
- Tuck pant legs into socks or boots and tuck loose shirts or jackets into pants.
- Wear light-colored clothing so that ticks can be spotted easily. If you see one, flick it off!
- Apply insect repellent containing Permethrin to pants, socks and shoes, and apply compounds containing DEET to exposed skin. Do not overuse these products.
- Walk in the center of trails to avoid overgrown grass and brush.
- After being outdoors in a tick-infested area, remove, wash and dry all clothing.
- Be sure to inspect your body thoroughly and carefully remove any attached ticks. To remove ticks attached to the skin, first pull them out gently with tweezers, taking care not to squeeze the tick's body. Then, apply an antiseptic to the bite site.



Pets should also be checked for ticks. A pet can also develop Lyme disease. Pets can carry ticks into the house. These ticks could fall off the animal and attach themselves to people inside the house, infecting them with their bite.

---

This document is for informational purposes and is not intended to be a substitute for the advice of a doctor or healthcare professional or a recommendation for any particular treatment plan. Like any printed material, it may become out of date over time. It is important that you rely on the advice of a doctor or a healthcare professional for your specific condition.

## Summary

Lyme disease is an infectious disease. Lyme disease is spread by a deer tick. If not treated, Lyme disease is a potentially serious disease that can lead to disabling arthritis and neurological problems. When a deer tick bites a person, the bacteria can transfer to the person's blood. The tick must stay attached to the person's skin for 2 days for infection to occur.

A person bitten by an infected deer tick develops a rash called erythema migrans or EM rash. It is usually the first symptom.

Over time, the bacteria travel through the blood stream and lymph nodes. They divide and multiply and produce harmful substances called toxins.

Lyme disease can also affect the nervous system, causing:

- Stiff neck.
- Severe headache.
- Temporary paralysis of facial muscles.
- Numbness, pain or weakness in the limbs.
- Poor motor coordination.

Most Lyme disease patients can be successfully treated with antibiotics. In general, the sooner such treatment is given, the quicker and more complete the recovery is. Sometimes it takes years for Lyme arthritis to fade away completely, but it usually goes away within a few weeks or months of antibiotic therapy.



Unfortunately, being infected with Lyme disease once does not guarantee that it will never happen again. Lyme disease can affect the same person more than once if he or she is re-infected with the Lyme disease bacterium.

---

This document is for informational purposes and is not intended to be a substitute for the advice of a doctor or healthcare professional or a recommendation for any particular treatment plan. Like any printed material, it may become out of date over time. It is important that you rely on the advice of a doctor or a healthcare professional for your specific condition.